



Melbourne Interventional Pulmonology Group Pty Ltd

Lung Function Request

Bookings: 9890 5200

Fax: 8372 0587

lungfunction@nickwilsmore.com.au

Patient Details

Name:

DOB: Contact No:

Address:

Referring Doctor

Name:

Address:

Provider No.: Date:

Signature:

Required Tests

- 1 Spirometry (pre +/- post) + FeNO (exhaled Nitric Oxide)
- 2 Spirometry (pre +/- Post) + TLCO (Gas Transfer) w/ Hb
- 3 Spirometry (pre +/- post) + Lung Volumes
- 4 Spirometry (pre +/- post) + CO
- 5 CO only (\$25 fee applies for single assessment)
- 6 Bronchial Provocation Testing (Mannitol - \$60 kit fee applies)
- 7 MIPS/MEPS/SNIPS

Please note: In order to bulk bill multiple appointments may be required.

Copy of Report to

Name:

Address:

Name:

Address:

Clinical Notes

Test on current inhaler medications Y / N

Appointment Details

Date: Time:

Location: Ekeru Building Level 3, Suite 1, 116 - 118 Thames St, Box Hill, 3128

Please see reverse for Patient Instructions.

Patient Instructions

PLEASE BRING THIS REFERRAL WITH YOU TO YOUR TEST

BEFORE ALL TESTS

We prefer you not to take any puffers (unless otherwise instructed)

- **NO** Ventolin, Respolin, Asmol, Bricanyl for 6 hours
- **NO** Atrovent, Seretide, Serevent, Symbicort, Oxis, Foradile, Spiriva or antihistamine tablets for 12 hours
- **NO** smoking within 4 hours

BEFORE BRONCHOPROVOCATIONS TESTS

As above, plus:

- **NO** tea, coffee, chocolate or caffeinated drinks to be consumed on the test day.
- **NO** oral steroids such as Prednisolone, Prednisone or Dexamethasone for 3 days - if advised/approved by your prescribing doctor.
- **NO** Antihistamines such as Dimetapp, Benadryl, Claratyne, Zyrtec or Telfast for 3 days.

BEFORE FeNO TESTS

As for "ALL TESTS" plus:

- **NO** food or drink for one hour before your test
- **AVOID** consuming any foods high in nitrogen on the day of the test including ham, bacon, salami, strawberries, currants, raspberries, cherries, gooseberries, spinach, lettuce, beetroot, celery, chervil radish and turnip tops.

IF YOU ARE MORE BREATHLESS THAN USUAL AFTER STOPPING THE ABOVE MEDICATIONS, PLEASE CONTACT YOUR DOCTOR AND RESUME TAKING THE MEDICATIONS.